

Quantum Gravity
from Integration
over Dirac ensembles

by John W. Barrett

Outline

I Spectral triples

- Commutative
- Non-commutative

II Quantum models

III Random Dirac ensembles

- Finite spectral triples
- dim 3 results

I: Commutative spectral triples

$(\mathcal{A}, \mathcal{H}, D)$

- \mathcal{A} : commutative $*$ -algebra
- \mathcal{H} : Hilbert space with action of \mathcal{A}
and commuting action of $\gamma, \gamma^2 = 1$
- D : $\mathcal{H} \rightarrow \mathcal{H}$
 $D\gamma = -\gamma D$ (s even)
 $\quad +\gamma D$ (s odd)
 $[[D, a], b] = 0$ for all $a, b \in \mathcal{A}$

Manifolds

$$(M, g_{\mu\nu}) \leftrightarrow (\mathcal{A}, \mathcal{H}, D)$$

- $\mathcal{A} = C^\infty(M, \mathbb{C})$
- $(*f)(x) = \bar{f}(x)$
- $\mathcal{H} = L^2(S, dV)$, S : bundle of spinors on M
- $\gamma =$ chirality of S
- $D = e_a^\mu(x) \gamma^a \nabla_\mu$, $e^2 = g$

Connes reconstruction: given dimension d ,
conditions on $(\mathcal{A}, \mathcal{H}, D)$ such that it is a d -manifold.

NC spectral triple

$(\mathcal{A}, \mathcal{H}, D)$

- \mathcal{A} : $*$ -algebra
- \mathcal{H} : Hilbert space, bimodule over \mathcal{A}
and commuting action of $\gamma, \gamma^2 = 1$
- D : $\mathcal{H} \rightarrow \mathcal{H}$
 $D\gamma = -\gamma D$ (s even)
 $\quad +\gamma D$ (s odd)
 $[[D, a \triangleright], \triangleleft b] = 0$ for all $a, b \in \mathcal{A}$

Real structure

$J: \mathcal{H} \rightarrow \mathcal{H}$, antilinear

- $J^2 = \pm 1$
- $JD = \pm' DJ$
- $J\gamma = \pm'' \gamma J$

Signs $\leftrightarrow s \in \mathbb{Z}/8$

Commutative case

$$Ja^*J^{-1} = a$$

M : spin manifold

Non-commutative case

$$\triangleleft a = J(a^* \triangleright)J^{-1}$$

SM internal space

$(\mathcal{A}_F, \mathcal{H}_F, D_F)$ finite real spectral triple, $s = 6$

- $\mathcal{A} = M_3(\mathbb{C}_{\mathbb{R}}) \oplus \mathbb{H} \oplus \mathbb{C}_{\mathbb{R}} = \{(m, q, \lambda)\}$
- $\mathcal{H} = \mathbb{C}^{96} = \langle l_L, e_R, \nu_R, q_L, d_R, u_R, \bar{l}_L, \bar{e}_R, \bar{\nu}_R, \bar{q}_L, \bar{d}_R, \bar{u}_R \rangle$
- $Jf = \bar{f}, J\bar{f} = f$

- $$D_F = \begin{pmatrix} 0 & M & G & 0 \\ M^* & 0 & 0 & H \\ G^* & 0 & 0 & \bar{M} \\ 0 & H^* & M^T & 0 \end{pmatrix}$$

basis $(f_L, f_R, \bar{f}_L, \bar{f}_R)$

particle	left action	right action	γ_F
l_L	q	λ	1
e_R	$\bar{\lambda}$	λ	-1
q_l	q	m^T	1
d_R	$\bar{\lambda}$	m^T	-1
u_R	λ	m^T	-1
\bar{l}_L	λ	q^T	-1
\bar{e}_R	λ	$\bar{\lambda}$	1
\bar{q}_l	m	q^T	-1
\bar{d}_R	m	$\bar{\lambda}$	1
\bar{u}_R	m	λ	1
ν_R	λ	λ	-1
$\bar{\nu}_R$	λ	λ	1

Vacuum of SM

$(\mathcal{A}_M, \mathcal{H}_M, D_M) = \text{spacetime}$

$(\mathcal{A}, \mathcal{H}, D_0) = (\mathcal{A}_M \otimes \mathcal{A}_F, \mathcal{H}_M \otimes \mathcal{H}_F, D_M \otimes 1 + \gamma_M \otimes D_F)$

D_0 is the vacuum of SM for the spacetime.

Physical fermion fields are in \mathcal{H}_+ : $\gamma_M \otimes \gamma_F = 1$

All bosonic fields: $D = D_0 + \sum_i a_i [D_0, b_i], \quad a_i, b_i \in \mathcal{A}$

II: Quantum models

Partition function for QG+SM:

$$Z(f) = \int e^{-S(D) + i \langle J\psi, D\psi \rangle} f(D, \psi) dD d\psi$$

$$D \in \mathcal{G}, \quad \psi \in \mathcal{H}_+$$

Issues:

- What is \mathcal{G} ?
- Is D_F fixed?
- What is S ?
- Are any axioms just e.o.m.?
- Functional integration?

III: Random Dirac models

Quantum models simplified:

- Assume fermions integrated already
- Fix \mathcal{H}, \mathcal{A} finite dimensional and NC
- $\mathcal{G} =$ all D satisfying real spectral triple axioms
- $S(D) = \text{tr } V(D)$, bounded below
- \int is ordinary integration on vector space \mathcal{G}

$$Z(f) = \int_{\mathcal{G}} e^{-S(D)} f(D) dD$$

Fuzzy spaces

$M(n) = n \times n$ matrices

$V =$ module for $\text{Cliff}(p,q)$

$s = q - p \pmod{8}$

- $\mathcal{A} = M(n, \mathbb{C}), M(n, \mathbb{R})$ or $M(n/2, \mathbb{H})$
- $\mathcal{H} = V \otimes M(n, \mathbb{C})$
- $\langle v \otimes m, v' \otimes m' \rangle = (v, v') \text{Tr } m^* m'$
- $\rho(a)(v \otimes m) = v \otimes (am)$
- $\Gamma(v \otimes m) = \gamma v \otimes m$
- $J(v \otimes m) = Cv \otimes m^*$

- Type (0,0)

$$D = 0$$

- Type (1,0)

$$D = \{H, \cdot\} + \gamma^1 \otimes \{H_1, \cdot\}$$

- Type (0,1)

$$D = [H, \cdot] + \gamma^1 \otimes [L_1, \cdot]$$

- Type (2,0)

$$D = \gamma^1 \otimes \{H_1, \cdot\} + \gamma^2 \otimes \{H_2, \cdot\}$$

- Type (1,1)

$$D = \gamma^1 \otimes \{H, \cdot\} + \gamma^2 \otimes [L, \cdot]$$

- Type (0,2)

$$D = \gamma^1 \otimes [L_1, \cdot] + \gamma^2 \otimes [L_2, \cdot]$$

Phase transition

$$S(D) = \text{tr } V(D)$$

$$V(D) = D^4 + g_2 D^2$$

$$S = \sum_{\lambda} \lambda^4 + g_2 \lambda^2$$





Figure 11: The potential $V = \lambda^4 + g_2 \lambda^2$ for $g_2 = -1, -1.5, -2, -2.5, -3, -3.5, -4, -4.5, -5$. The lines are coloured from red ($g_2 = -1$) through to yellow ($g_2 = -5$).



(c) Type (2, 0)

Monte Carlo
Eigenvalue distribution

JWB + L. Glaser 2016

Numerical simulation of random Dirac operators

Thesis submitted to the University of Nottingham for the degree of
Doctor of Philosophy, March 2022.

Mauro D'Arcangelo
14302771

Supervised by

John W. Barrett
Sven Gnutzmann

Decompose $m_\mu = t_\mu 1 + v_\mu$
with $\text{tr } v_\mu = 0$



Type $(p, q) = (3, 0)$:

$$D = 1 \otimes [m_0, \cdot] + \sum_1^3 \sigma_i \otimes \{m_i, \cdot\}$$

Type $(p, q) = (0, 3)$:

$$D = 1 \otimes \{m_0, \cdot\} + \sum_1^3 \sigma_i \otimes [m_i, \cdot]$$

Type (3,0)



$$D = 1 \otimes [m_0, \cdot] + \sum_1^3 \sigma_i \otimes \{m_i, \cdot\}$$





Figure 6.8: Monte Carlo history of t_1 , t_2 and t_3 in region II of the (3,0) model at $g_2 = -6$, $n = 8$. The solid orange sphere is a guide for the eyes.

2nd order transition to commutative phase

Type (0,3)



$$D = 1 \otimes \{m_0, \cdot\} + \sum_1^3 \sigma_a \otimes [m_a, \cdot]$$

g_2	Chain 1	Chain 2	Chain 3	Chain 4	Fuzzy sphere	$-g_2/16$
-300	18.6946(3)	18.6946(2)	18.6945(2)	18.6946(2)	18.6951	18.75
-150	9.3465(3)	9.3740(3)	9.3465(2)	9.3739(2)	9.3476	9.375
-100	6.2301(2)	6.2301(3)	6.2301(2)	6.2301(3)	6.2317	6.25

Fuzzy sphere $v_a = Rl_a$, $a = 1,2,3$

$$[l_a, l_b] = \sum_c i\epsilon_{abc} l_c, \text{ irreducible}$$

$$\frac{1}{n} \text{Tr} v_c^2 = -\frac{g_2}{8} \frac{n^2 - 1}{2n^2 - 1} \approx -\frac{g_2}{16}, \quad c = 1, 2, 3$$




Figure 6.6: Model (0,3), eigenvalue density of v_1/R (purple) and $i[v_2, v_3]/R^2$ (green) for $n = 12$, $g_2 = -300$. The spectrum is compatible with an $su(2)$ solution.

Conclusion

- Would like to model (Euclidean) quantum spacetime with a random Dirac model.
- This supposes spacetime has some NC structure. If it does, there is a good explanation of the Planck scale.
- Understanding the vacuum in such models is crucial to explaining the physical picture.

References

Real spectral triples

A. Connes. Noncommutative geometry and reality. 1995.

JWB, Introduction to Non-commutative Geometry. 2022 <https://johnwbarrett.wordpress.com>

Standard Model

JWB, A Lorentzian version of the non-commutative geometry of the standard model of particle physics. 2006

A. Connes. Noncommutative geometry and the standard model with neutrino mixing. 2006

Random spectral triples

JWB, L. Glaser, Monte Carlo simulations of random non-commutative geometries. 2016

M. D'Arcangelo, Numerical simulation of random Dirac operators. PhD thesis, University of Nottingham. 2022 <https://eprints.nottingham.ac.uk/theses/>

H. Hessam, M. Khalkhali, N. Pagliaroli, and L. Verhoeven, From Noncommutative Geometry to Random Matrix Theory 2022